

## LY75 Antibody (Center) Blocking Peptide

Synthetic peptide Catalog # BP8994c

## **Specification**

## LY75 Antibody (Center) Blocking Peptide - Product Information

**Primary Accession** 

060449

# LY75 Antibody (Center) Blocking Peptide - Additional Information

Gene ID 100526664;4065

#### **Other Names**

Lymphocyte antigen 75, Ly-75, C-type lectin domain family 13 member B, DEC-205, gp200-MR6, CD205, LY75, CD205, CLEC13B

## **Target/Specificity**

The synthetic peptide sequence used to generate the antibody <a

href=/products/AP8994c>AP8994c</a> was selected from the Center region of human LY75. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## LY75 Antibody (Center) Blocking Peptide - Protein Information

Name LY75

Synonyms CD205, CLEC13B

#### **Function**

Acts as an endocytic receptor to direct captured antigens from the extracellular space to a specialized antigen-processing compartment (By similarity). Causes reduced proliferation of B-lymphocytes.

#### **Cellular Location**

Membrane; Single-pass type I membrane protein

## **Tissue Location**

Expressed in spleen, thymus, colon and peripheral blood lymphocytes. Detected in myeloid and



B-lymphoid cell lines Isoform 2 and isoform 3 are expressed in malignant Hodgkin lymphoma cells called Hodgkin and Reed-Sternberg (HRS) cells

# LY75 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

#### • Blocking Peptides

LY75 Antibody (Center) Blocking Peptide - Images

# LY75 Antibody (Center) Blocking Peptide - Background

LY75 acts as an endocytic receptor to direct captured antigens from the extracellular space to a specialized antigen-processing compartment. It causes reduced proliferation of B-lymphocytes.

# LY75 Antibody (Center) Blocking Peptide - References

Kato, M., et.al., Immunogenetics 47 (6), 442-450 (1998) McKay, P.F., et.al., Eur. J. Immunol. 28 (12), 4071-4083 (1998)