

ND3 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP9011b**Specification**

ND3 Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession [P03897](#)**ND3 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 4537**Other Names**

NADH-ubiquinone oxidoreductase chain 3, NADH dehydrogenase subunit 3, MT-ND3, MTND3, NADH3, ND3

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP9011b](/products/AP9011b) was selected from the C-term region of human ND3. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ND3 Antibody (C-term) Blocking Peptide - Protein Information**Name** MT-ND3 ([HGNC:7458](#))**Synonyms** MTND3, NADH3, ND3**Function**

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which catalyzes electron transfer from NADH through the respiratory chain, using ubiquinone as an electron acceptor (PubMed: <http://www.uniprot.org/citations/25118196> target="_blank">25118196). Essential for the catalytic activity of complex I (PubMed: <http://www.uniprot.org/citations/25118196> target="_blank">25118196).

Cellular Location

Mitochondrion inner membrane {ECO:0000250|UniProtKB:P03898}; Multi-pass membrane protein

ND3 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

ND3 Antibody (C-term) Blocking Peptide - Images

ND3 Antibody (C-term) Blocking Peptide - Background

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone (By similarity).

ND3 Antibody (C-term) Blocking Peptide - References

Sarzi E., et.al., Am. J. Med. Genet. A 143:33-41(2007). Taylor R.W., et.al., Ann. Neurol. 50:104-107(2001).