

DSTN Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP9189c**Specification**

DSTN Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [P60981](#)**DSTN Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 11034**Other Names**

Destrin, Actin-depolymerizing factor, ADF, DSTN, ACTDP, DSN

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP9189c](/products/AP9189c) was selected from the Center region of human DSTN. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

DSTN Antibody (Center) Blocking Peptide - Protein Information**Name** DSTN**Synonyms** ACTDP, DSN**Function**

Actin-depolymerizing protein. Severs actin filaments (F- actin) and binds to actin monomers (G-actin). Acts in a pH-independent manner.

Tissue Location

Widely distributed in various tissues.

DSTN Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

DSTN Antibody (Center) Blocking Peptide - Images

DSTN Antibody (Center) Blocking Peptide - Background

DSTN belongs to the actin-binding proteins ADF family. This family of proteins is responsible for enhancing the turnover rate of actin in vivo. This protein encodes the actin depolymerizing protein that severs actin filaments (F-actin) and binds to actin monomers (G-actin).

DSTN Antibody (Center) Blocking Peptide - References

Estornes,Y., et.al., Int. J. Cancer 121 (10), 2162-2171 (2007)Mannherz,H.G., et.al., J. Mol. Biol. 366 (3), 745-755 (2007)