

TYW2 Antibody (C-term) Blocking Peptide

Synthetic peptide Catalog # BP9305b

Specification

TYW2 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

Q53H54

TYW2 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 55039

Other Names

tRNA wybutosine-synthesizing protein 2 homolog, tRNA-yW-synthesizing protein 2, tRNA(Phe) (4-demethylwyosine(37)-C(7)) aminocarboxypropyltransferase, TRMT12, TRM12, TYW2

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP9305b was selected from the C-term region of human TYW2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

TYW2 Antibody (C-term) Blocking Peptide - Protein Information


Name TRMT12

Synonyms TRM12, TYW2

Function

S-adenosyl-L-methionine-dependent transferase that acts as a component of the wybutosine biosynthesis pathway. Wybutosine is a hyper modified guanosine with a tricyclic base found at the 3'-position adjacent to the anticodon of eukaryotic phenylalanine tRNA. Catalyzes the transfer of the alpha-amino-alpha-carboxypropyl (acp) group from S- adenosyl-L-methionine to the C-7 position of 4-demethylwyosine (imG-14) to produce wybutosine-86.

TYW2 Antibody (C-term) Blocking Peptide - Protocols





Tel: 858.875.1900 Fax: 858.875.1999

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

TYW2 Antibody (C-term) Blocking Peptide - Images

TYW2 Antibody (C-term) Blocking Peptide - Background

TYW2 is a hypermodified guanosine at the 3-prime position adjacent to the anticodon of phenylalanine tRNA that stabilizes codon-anticodon interactions during decoding on the ribosome. TRMT12 is the human homolog of a yeast gene essential for yW synthesis.

TYW2 Antibody (C-term) Blocking Peptide - References

Rodriguez, V., et.al., Genes Chromosomes Cancer 46 (7), 694-707 (2007) Noma, A. et.al., Nucleic Acids Symp Ser (Oxf) 50, 65-66 (2006)Kalhor, H.R., et.al., Biochem. Biophys. Res. Commun. 334 (2), 433-440 (2005)