

H2A1J Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP9426b**Specification**

H2A1J Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession [Q99878](#)**H2A1J Antibody (C-term) Blocking Peptide - Additional Information**

Gene ID 8331

Other Names

Histone H2A type 1-J, Histone H2A/e, HIST1H2AJ, H2AFE

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

H2A1J Antibody (C-term) Blocking Peptide - Protein InformationName H2AC14 ([HGNC:4727](#))**Function**

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location

Nucleus. Chromosome.

H2A1J Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

H2A1J Antibody (C-term) Blocking Peptide - Images

H2A1J Antibody (C-term) Blocking Peptide - Background

H2A1J are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H2A family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3.

H2A1J Antibody (C-term) Blocking Peptide - References

Lusic, M., et al. EMBO J. 22(24):6550-6561(2003)Marzluff, W.F., et al. Genomics 80(5):487-498(2002)Andersen, J.S., et al. Curr. Biol. 12(1):1-11(2002)Deng, L., et al. Virology 289(2):312-326(2001)