

SGMS2 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP9801b**Specification**

SGMS2 Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession [Q8NHU3](#)**SGMS2 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 166929**Other Names**

Phosphatidylcholine:ceramide cholinephosphotransferase 2, Sphingomyelin synthase 2, SGMS2, SMS2

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

SGMS2 Antibody (C-term) Blocking Peptide - Protein Information**Name** SGMS2 {ECO:0000303|PubMed:30779713, ECO:0000312|HGNC:HGNC:28395}**Function**

Sphingomyelin synthase that primarily contributes to sphingomyelin synthesis and homeostasis at the plasma membrane. Catalyzes the reversible transfer of phosphocholine moiety in sphingomyelin biosynthesis: in the forward reaction transfers phosphocholine head group of phosphatidylcholine (PC) on to ceramide (CER) to form ceramide phosphocholine (sphingomyelin, SM) and diacylglycerol (DAG) as by-product, and in the reverse reaction transfers phosphocholine from SM to DAG to form PC and CER. The direction of the reaction appears to depend on the levels of CER and DAG in the plasma membrane (PubMed:14685263, PubMed:17449912, PubMed:17982138, PubMed:18370930). Does not use free phosphorylcholine or CDP-choline as donors (PubMed:14685263). Can also transfer phosphoethanolamine head group of phosphatidylethanolamine (PE) on to ceramide (CER) to form ceramide phosphoethanolamine (CPE) (PubMed:19454763). Regulates receptor-mediated signal transduction via mitogenic DAG and proapoptotic CER, as well as via SM,

a structural component of membrane rafts that serve as platforms for signal transduction and protein sorting (PubMed:17449912, PubMed:17982138). To a lesser extent, plays a role in secretory transport via regulation of DAG pool at the Golgi apparatus and its downstream effects on PRKD1 (PubMed:18370930, PubMed:21980337). Required for normal bone matrix mineralization (PubMed:30779713).

Cellular Location

Cell membrane; Multi-pass membrane protein. Golgi apparatus membrane; Multi-pass membrane protein. Note=Primarily localized at the plasma membrane with a small fraction at the Golgi apparatus.

Tissue Location

Brain, heart, kidney, liver, muscle and stomach. Also expressed in a number of cell lines such as carcinoma HeLa cells, hepatoma Hep-G2 cells, and colon carcinoma Caco-2 cells

SGMS2 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

SGMS2 Antibody (C-term) Blocking Peptide - Images

SGMS2 Antibody (C-term) Blocking Peptide - Background

Sphingomyelin, a major component of cell and Golgi membranes, is made by the transfer of phosphocholine from phosphatidylcholine onto ceramide, with diacylglycerol as a side product. The protein encoded by this gene is an enzyme that catalyzes this reaction primarily at the cell membrane. The synthesis is reversible, and this enzyme can catalyze the reaction in either direction. The encoded protein is required for cell growth. Three transcript variants encoding the same protein have been found for this gene.

SGMS2 Antibody (C-term) Blocking Peptide - References

Ternes, P., et al. J. Lipid Res. 50(11):2270-2277(2009)Liu, J., et al. Arterioscler. Thromb. Vasc. Biol. 29(6):850-856(2009)Tani, M., et al. Biochem. Biophys. Res. Commun. 381(3):328-332(2009)Wang, W., et al. Plant Cell 20(11):3163-3179(2008)Yeang, C., et al. Biochim. Biophys. Acta 1781(10):610-617(2008)Villani, M., et al. Biochem. J. 414(1):31-41(2008)