

**Recombinant Human BAFF**  
**Catalog # PBG10025****Specification**

---

**Recombinant Human BAFF - Product Information****Recombinant Human BAFF - Additional Information****Description**

BAFF, a member of the TNF family of ligands, is expressed in T cells, macrophages, monocytes and dendritic cells. BAFF is involved in stimulation of B and T cell function, and is an important survival and maturation factor for peripheral B cells. BAFF signals through three different TNF receptors TACI, BCMA and BAFF-R. The human BAFF gene codes for a 285 amino acid type II transmembrane protein containing a 46 amino acid cytoplasmic domain, a 21 amino acid transmembrane domain, and a 218 amino acid extracellular domain. Recombinant human soluble BAFF is a 152 amino acid polypeptide (17.0 kDa), which contains the TNF-like portion of the extracellular domain of BAFF.

**Biological Activity**

Determined by a M splenocyte survival assay. The expected  $ED_{50}$  for this effect is 0.5-2.0 µg/ml.

**Authenticity**

Verified by N-terminal and Mass Spectrometry analyses (when applicable).

**Endotoxin**

Endotoxin level is <0.1 ng/ µg of protein (<1EU/ µg).

**Protein Content**

Verified by UV Spectroscopy and/or SDS-PAGE gel.

**Storage**

-20°C

**Precautions**

Recombinant Human BAFF is for research use only and not for use in diagnostic or therapeutic procedures.

**Recombinant Human BAFF - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

## **Recombinant Human BAFF - Images**