

Recombinant Human FGF-17
Catalog # PBG10093**Specification**

Recombinant Human FGF-17 - Product Information**Recombinant Human FGF-17 - Additional Information****Description**

FGF-17 is a heparin binding growth factor that is a member of the FGF family. Proteins of this family play a central role during prenatal development and postnatal growth and regeneration of a variety of tissues, by promoting cellular proliferation and differentiation. FGF-17 signals through the FGFR 1c, 2c, 3c, and 4. FGF-17 signals induction and patterning of embryonic brain. Recombinant human FGF-17 is a 22.7 kDa protein consisting of 195 amino acid residues.

Biological Activity

Assay #1: The **ED₅₀** as determined by the dose-dependent stimulation of thymidine uptake by BaF3 cells expressing FGF-receptors is ≤ 0.5 ng/ml, corresponding to a specific activity of $\geq 2 \times 10^6$ units/mg.
Assay #2: The **ED₅₀** was determined by a cell proliferation assay using balb/c 3T3 cells is ≤ 10.0 ng/ml, corresponding to a specific activity of $\geq 1 \times 10^5$ units/mg.

Authenticity

Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Endotoxin

Endotoxin level is <0.1 ng/ μ g of protein (<1 EU/ μ g).

Protein Content

Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage

-20°C

Precautions

Recombinant Human FGF-17 is for research use only and not for use in diagnostic or therapeutic procedures.

Recombinant Human FGF-17 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Recombinant Human FGF-17 - Images