

Recombinant Human FGF-acidic

Catalog # PBG10105

Specification

Recombinant Human FGF-acidic - Product Information

Recombinant Human FGF-acidic - Additional Information

Description

FGF-acidic is one of 23 known members of the FGF family. Proteins of this family play a central role during prenatal development and postnatal growth and regeneration of a variety of tissues, by promoting cellular proliferation and differentiation. FGF-acidic is a non-glycosylated heparin binding growth factor that is expressed in the brain, kidney, retina, smooth muscle cells, bone matrix, osteoblasts, astrocytes and endothelial cells. FGF-acidic has the ability to signal through all the FGF receptors. Recombinant human FGF-acidic is a 16.0 kDa protein consisting of 141 amino acid residues.

BiologicalActivity

Assay #1: The ED₅₀ as determined by the dose-dependent stimulation of thymidine uptake by BaF3 cells expressing FGF receptors is ≤ 10 ng/ml, corresponding to a specific activity of $\geq 1 \times 10$ ⁵ units/mg.
br /> Assay #2: The ED₅₀ was determined by a cell proliferation assay using balb/c 3T3 cells is ≤ 0.5 ng/ml in the presence of 10 µg/ml heparin, corresponding to a specific activity of $\geq 2 \times 10$ ⁶ units/mg.

Authenticity Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Endotoxin Endotoxin level is <0.1 ng/ μ g of protein (<1EU/ μ g).

Protein Content Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage -20°C

Precautions

Recombinant Human FGF-acidic is for research use only and not for use in diagnostic or therapeutic procedures.

Recombinant Human FGF-acidic - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot



- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Recombinant Human FGF-acidic - Images