10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Recombinant Human GDF-3
Catalog # PBG10134

Specification

Recombinant Human GDF-3 - Product Information

Recombinant Human GDF-3 - Additional Information

Description
GDF-3 is a member of the TGF-β superfamily of growth and differentiation factors, and is highly homologous to GDF-9. Unlike most TGF-β family members, GDF-3 and GDF-9 are not disulfide-linked dimers. GDF-3 is expressed in adult bone marrow, spleen, thymus, and adipose tissue. The expression of GDF-3 is upregulated in high-fat-fed wild-type FABP4/aP2 null mice and was associated with obesity, but not with the related hyperglycemia/hyperinsulinemia which characterizes Type 2 diabetes. Recombinant human GDF-3 is a 26.0 kDa non-disulfide-linked homodimer containing two 114 amino acid polypeptide chains.

BiologicalActivity
Determined by its ability to inhibit induced alkaline phosphatase production by ATDC-5 chondrogenic cells. The ED₅₀ for this effect is 100-150 ng/ml.

Authenticity
Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Endotoxin
Endotoxin level is <0.1 ng/ µg of protein (<1EU/ µg).

Protein Content
Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage
-20°C

Precautions
Recombinant Human GDF-3 is for research use only and not for use in diagnostic or therapeutic procedures.

Recombinant Human GDF-3 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety

- Cell Culture

Recombinant Human GDF-3 - Images