

**Recombinant Human ICAM-1**  
**Catalog # PBG10161****Specification**

---

**Recombinant Human ICAM-1 - Product Information****Recombinant Human ICAM-1 - Additional Information****Description**

ICAMs are members of the Ig superfamily of calcium-independent transmembrane glycoproteins. ICAM-1 is a ligand for lymphocyte function-associated (LFA) and Mac-1 integrins and the major human rhinovirus receptor. The primary function of ICAM-1 is to provide adhesion between endothelial cells and leukocytes after stress or injury. The human ICAM-1 gene codes for a 505 amino acid transmembrane glycoprotein containing a 29 amino acid cytoplasmic domain, a 23 amino acid transmembrane domain, and a 453 amino acid extracellular domain. Recombinant human ICAM-1 is a 49.5 kDa glycoprotein comprising the extracellular domain (453 amino acid residues) of ICAM-1. Monomeric glycosylated ICAM-1 migrates at an apparent molecular weight of approximately 72.0-80.0 kDa by SDS-PAGE analysis under reducing conditions.

**Biological Activity**

ICAM-1 supported the adhesion of more than 50% PMA-stimulated HSB2 cells to plates coated at 25µg/ml ICAM-1 concentration.

**Authenticity**

Verified by N-terminal and Mass Spectrometry analyses (when applicable).

**Endotoxin**

Endotoxin level is <0.1 ng/ µg of protein (<1EU/ µg).

**Protein Content**

Verified by UV Spectroscopy and/or SDS-PAGE gel.

**Storage**

-20°C

**Precautions**

Recombinant Human ICAM-1 is for research use only and not for use in diagnostic or therapeutic procedures.

**Recombinant Human ICAM-1 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Recombinant Human ICAM-1 - Images**