

# Recombinant Murine IL-1β

Catalog # PBG10212

## Specification

# **Recombinant Murine IL-1**<sup>β</sup> - **Product Information**

## Recombinant Murine IL-1β - Additional Information

### Description

IL-1 $\beta$  is a proinflammatory cytokine produced in a variety of cells including monocytes, tissue macrophages, keratinocytes and other epithelial cells. Both IL-1 $\alpha$  and IL-1 $\beta$  binds to the same receptor and has similar if not identical biological properties. These cytokines have a broad range of activities including, stimulation of thymocyte proliferation, by inducing IL-2 release, B-cell maturation and proliferation, mitogenic FGF-like activity and the ability to stimulate the release of prostaglandin and collagenase from synovial cells. However, whereas IL-1 $\beta$  is a secreted cytokine, IL-1 $\alpha$  is predominantly a cell-associated cytokine. Recombinant murine IL-1 $\beta$  is a 17.5 kDa protein containing 153 amino acid residues.

#### BiologicalActivity

The ED<sub>50</sub> as determined by the dose-dependent stimulation of murine D10S cells is  $\leq 0.002$  ng/ml, corresponding to a specific activity of  $\geq 5 \times 10$ <sup>8</sup> units/mg.

Authenticity Verified by N-terminal and Mass Spectrometry analyses (when applicable).

**Endotoxin** Endotoxin level is <0.1 ng/ μg of protein (<1EU/ μg).

### **Protein Content**

Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage -20°C

Precautions

Recombinant Murine IL-1 $\beta$  is for research use only and not for use in diagnostic or therapeutic procedures.

### **Recombinant Murine IL-1β - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation



<u>Flow Cytomety</u>
<u>Cell Culture</u> **Recombinant Murine IL-1β - Images**