

Recombinant Murine SDF-1α (CXCL12)

Catalog # PBG10404

Specification

Recombinant Murine SDF-1α (CXCL12) - Product Information

Recombinant Murine SDF-1α (CXCL12) - Additional Information

Description

SDF- 1α and β are stromal derived CXC chemokines, and signal through the CXCR4 receptor. SDF- 1α and β chemoattract B and T cells, and have been shown to induce migration of CD34+ stem cells. Additionally, the SDF-1 proteins exert HIV suppressive activity in cells expressing the CXCR4 receptor. Recombinant murine SDF- 1α is a 7.9 kDa protein containing 68 amino acid residues.

BiologicalActivity

Determined by its ability to chemoattract human peripheral blood monocytes using a concentration of 50-100 ng/ml.

Authenticity

Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Endotoxin

Endotoxin level is <0.1 ng/ μg of protein ($<1EU/ \mu g$).

Protein Content

Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage

-20°C

Precautions

Recombinant Murine SDF- 1α (CXCL12) is for research use only and not for use in diagnostic or therapeutic procedures.

Recombinant Murine SDF-1α (CXCL12) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Recombinant Murine SDF-1α (CXCL12) - Images



