

Jagged 1 Blocking Peptide
Catalog # PBV10047b**Specification**

Jagged 1 Blocking Peptide - Product Information

Primary Accession	Q63722
Gene ID	29146
Calculated MW	134326

Jagged 1 Blocking Peptide - Additional Information**Gene ID** 29146**Application & Usage**

The peptide is used for blocking the antibody activity of Jagged 1. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.

Other Names

Protein jagged-1, Jagged1, CD339, Jag1

Target/Specificity

Jagged 1

Formulation

50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

Jagged 1 Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

Jagged 1 Blocking Peptide - Protein Information**Name** Jag1**Function**

Ligand for multiple Notch receptors and involved in the mediation of Notch signaling. May be involved in cell-fate decisions during hematopoiesis. Enhances fibroblast growth factor-induced angiogenesis (in vitro). Seems to be involved in early and late stages of mammalian cardiovascular development. Inhibits myoblast differentiation. May regulate fibroblast growth

factor-induced angiogenesis.

Cellular Location

Membrane; Single-pass type I membrane protein. Cell membrane
{ECO:0000250|UniProtKB:P78504}

Tissue Location

Widely expressed in a variety of tissues.

Jagged 1 Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Jagged 1 Blocking Peptide - Images