

TRIM71 Blocking Peptide

Catalog # PBV10096b

Specification

TRIM71 Blocking Peptide - Product Information

Primary Accession
Gene ID
Calculated MW
Q201W2
131405
93385

TRIM71 Blocking Peptide - Additional Information

Gene ID 131405

Application & Usage The peptide is used for blocking the

antibody activity of TRIM71. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for

30-60 minutes at 37°C.

Other Names

E3 ubiquitin-protein ligase TRIM71, 6.3.2.-, Protein lin-41 homolog, Tripartite motif-containing protein 71, TRIM71, LIN41

Target/Specificity

TRIM71

Formulation

 $50~\mu g$ (0.2 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 0.1% BSA and 0.02% thimerosal.

Reconstitution & Storage

-20 °C

Background Descriptions

Precautions

TRIM71 Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

TRIM71 Blocking Peptide - Protein Information

Name TRIM71 (<u>HGNC:32669</u>)

Synonyms LIN41

Function

E3 ubiquitin-protein ligase that cooperates with the microRNAs (miRNAs) machinery and promotes



embryonic stem cells proliferation and maintenance (Probable). Binds to miRNAs and associates with AGO2, participating in post-transcriptional repression of transcripts such as CDKN1A (By similarity). In addition, participates in post-transcriptional mRNA repression in a miRNA independent mechanism (PubMed:23125361). Facilitates the G1-S transition to promote rapid embryonic stem cell self-renewal by repressing CDKN1A expression. Required to maintain proliferation and prevent premature differentiation of neural progenitor cells during early neural development: positively regulates FGF signaling by controlling the stability of SHCBP1 (By similarity). Specific regulator of miRNA biogenesis. Binds to miRNA MIR29A hairpin and postranscriptionally modulates MIR29A levels, which indirectly regulates TET proteins expression (PubMed:28431233,a>).

Cellular Location Cytoplasm, P-body

Tissue LocationSpecifically expressed in testis.

TRIM71 Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

TRIM71 Blocking Peptide - Images