

C-Peptide Blocking Peptide
Catalog # PBV10118b**Specification**

C-Peptide Blocking Peptide - Product Information**C-Peptide Blocking Peptide - Additional Information****Application & Usage**

The peptide is used for blocking the antibody activity of C-peptide. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.

Target/Specificity

C-Peptide

Formulation

50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

C-Peptide Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

C-Peptide Blocking Peptide - Protein Information**C-Peptide Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C-Peptide Blocking Peptide - Images

