10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Cytochrome c Blocking Peptide

Catalog # PBV10143b

Specification

Cytochrome c Blocking Peptide - Product Information

Cytochrome c Blocking Peptide - Additional Information

Application & Usage

The peptide is used for blocking the antibody activity of active r-Cytochrome c. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30 minutes at 37°C

Target/Specificity
Cytochrome c

Formulation
50 µg (0.2 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 0.1% BSA and 0.02% thimerosal.

Reconstitution & Storage
-20 °C

Background Descriptions

Precautions
Cytochrome c Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

Cytochrome c Blocking Peptide - Protein Information

Cytochrome c Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Cytochrome c Blocking Peptide - Images

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999