

**Stat3 Blocking Peptide**  
**Catalog # PBV10198b****Specification**

---

**Stat3 Blocking Peptide - Product Information**

Primary Accession	<a href="#">P42227</a>
Other Accession	<a href="#">AAA19452.1</a>
Gene ID	<b>20848</b>
Calculated MW	<b>88054</b>

**Stat3 Blocking Peptide - Additional Information****Gene ID** 20848**Application & Usage**

The peptide is used for blocking the antibody activity of Stat3. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.

**Other Names**

Signal transducer and activator of transcription 3, Acute-phase response factor, Stat3, Aprf

**Target/Specificity**

Stat3

**Formulation**

50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

**Reconstitution & Storage**

-20 °C

**Background Descriptions****Precautions**

Stat3 Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

**Stat3 Blocking Peptide - Protein Information****Name** Stat3 {ECO:0000312|MGI:MGI:103038}**Function**

Signal transducer and transcription activator that mediates cellular responses to interleukins, KITLG/SCF, LEP and other growth factors (By similarity). Once activated, recruits coactivators, such as NCOA1 or MED1, to the promoter region of the target gene (By similarity). May mediate cellular

responses to activated FGFR1, FGFR2, FGFR3 and FGFR4 (By similarity). Upon activation of IL6ST/gp130 signaling by interleukin-6 (IL6), binds to the IL6-responsive elements identified in the promoters of various acute-phase protein genes (By similarity). Activated by IL31 through IL31RA (By similarity). Acts as a regulator of inflammatory response by regulating differentiation of naive CD4(+) T-cells into T-helper Th17 or regulatory T-cells (Treg): acetylation promotes its transcription activity and cell differentiation while deacetylation and oxidation of lysine residues by LOXL3 inhibits differentiation (By similarity). Involved in cell cycle regulation by inducing the expression of key genes for the progression from G1 to S phase, such as CCND1 (By similarity). Mediates the effects of LEP on melanocortin production, body energy homeostasis and lactation (PubMed:<a href="http://www.uniprot.org/citations/12594516" target="\_blank">12594516</a>). May play an apoptotic role by transactivating BIRC5 expression under LEP activation (PubMed:<a href="http://www.uniprot.org/citations/16825198" target="\_blank">16825198</a>). Cytoplasmic STAT3 represses macroautophagy by inhibiting EIF2AK2/PKR activity (By similarity). Plays a crucial role in basal beta cell functions, such as regulation of insulin secretion (PubMed:<a href="http://www.uniprot.org/citations/20215569" target="\_blank">20215569</a>). Following JAK/STAT signaling activation and as part of a complex with NFATC3 and NFATC4, binds to the alpha-beta E4 promoter region of CRYAB and activates transcription in cardiomyocytes (PubMed:<a href="http://www.uniprot.org/citations/19538478" target="\_blank">19538478</a>). Plays an important role in host defense in methicillin-resistant S.aureus lung infection by regulating the expression of the antimicrobial lectin REG3G (PubMed:<a href="http://www.uniprot.org/citations/23401489" target="\_blank">23401489</a>).

#### **Cellular Location**

Cytoplasm {ECO:0000250|UniProtKB:P40763}. Nucleus. Note=Predominantly present in the cytoplasm without stimuli. Upon leukemia inhibitory factor (LIF) stimulation, accumulates in the nucleus. The complex composed of BART and ARL2 plays an important role in the nuclear translocation and retention of STAT3 (By similarity). Shuttles between the nucleus and the cytoplasm. Translocated into the nucleus upon tyrosine phosphorylation and dimerization, in response to signaling by activated FGFR1, FGFR2, FGFR3 or FGFR4. Constitutive nuclear presence is independent of tyrosine phosphorylation. Translocates to the nucleus in the presence of EDN1 (By similarity). {ECO:0000250, ECO:0000250|UniProtKB:P52631}

#### **Tissue Location**

Expressed in ventricular cardiomyocytes (at protein level) (PubMed:19538478). Expressed in the lung (at protein level) (PubMed:23401489). Expressed in the liver, spleen and kidney (PubMed:7512451).

### **Stat3 Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Stat3 Blocking Peptide - Images**