10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

BD-1, human recombinant protein

β-defensin 1, BD-1, Defensin β 1, hBD-1, HBD1, HBP1, DEFB1, HBD-1, HBP-1, DEFB101, DEFB-1, MGC51822

Catalog # PBV10289r

Specification

BD-1, human recombinant protein - Product info

Primary Accession	P60022
Calculated MW	**5.0 kDa KDa**

BD-1, human recombinant protein - Additional Info

Gene ID	**1672**
Gene Symbol	**DEFB1**

Other Names
β-defensin 1, BD-1, Defensin β 1, hBD-1, HBD1, HBP1, DEFB1, HBD-1, HBP-1, DEFB101, DEFB-1, MGC51822

Gene Source	**Human**
Source	**E. coli**
Assay&Purity	**SDS-PAGE; ≥98%**
Assay2&Purity2	**HPLC;**
Recombinant	**Yes**
Results	**0.1-1 µg/ml**

Target/Specificity
BD-1

Application Notes
Reconstitute in dH_2O to a concentration of 0.1-1.0 mg/ml. This solution can then be diluted into other aqueous buffers and stored at 4°C for 1 week or –20°C for future use.

Format
Lyophilized protein

Storage
-20°C; Lyophilized from a concentrated (1 mg/ml) solution containing 20 mM PBS pH-7.4 and 130 mM sodium chloride

BD-1, human recombinant protein - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation

- Flow Cytomety
- Cell Culture

BD-1, human recombinant protein - Images

BD-1, human recombinant protein - Background

The Defensin family are highly similar in their protein sequence and are microbicidal & cytotoxic peptides made by neutrophils. β Defensin-1 (BD-1) is an antimicrobial peptide having the resistance of epithelial surfaces to microbial colonization. BD-1 has close proximity to Defensin α-1 and has been implicated in the pathogenesis of cystic fibrosis. Skin of patients having atopic dermatitis patients and mycosis fungoides (non-lesional and lesional) show lower human BD-1 mRNA expression and higher human BD-2 and human BD-3 mRNA expression. β Defensin is highly expressed by epithelial cells. BD-1 may play a role in the pathogenesis of severe sepsis. Variation in human BD-1 contributes to asthma diagnosis, with apparent gender-specific effects. Human BD-3 is a dimer, while Human BD-1 and Human BD-2 are monomeric. Recombinant human β Defensin-1 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 47 amino acids and having a molecular mass of 5 kDa.

BD-1, human recombinant protein - References

Liu L.,et al.Genomics 43:316-320(1997).
Zhao C.,et al.Submitted (OCT-1996) to the EMBL/GenBank/DDBJ databases.
McCray P.B. Jr.,et al.Am. J. Respir. Cell Mol. Biol. 16:343-349(1997).
Bensch K.W.,et al.FEBS Lett. 368:331-335(1995).
Kluever E.,et al.J. Pept. Res. 59:241-248(2002).