

Tubulin Blocking Peptide
Catalog # PBV10332b**Specification**

Tubulin Blocking Peptide - Product Information

Other Accession [AAP36356.1](#)
Gene ID **30584215**

Tubulin Blocking Peptide - Additional Information**Application & Usage**

The peptide is used for blocking the antibody activity of Tubulin. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.

Target/Specificity

Tubulin

Formulation

50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

Tubulin Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

Tubulin Blocking Peptide - Protein Information**Tubulin Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

Tubulin Blocking Peptide - Images