

## **Calnexin Blocking Peptide**

Catalog # PBV10362b

## **Specification**

## **Calnexin Blocking Peptide - Product Information**

Primary Accession P27824
Gene ID 821
Calculated MW 67568

## **Calnexin Blocking Peptide - Additional Information**

Gene ID 821

Application & Usage The peptide is used for blocking the

antibody activity of Calnexin. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for

30-60 minutes at 37°C.

**Other Names** 

Calnexin, IP90, Major histocompatibility complex class I antigen-binding protein p88, p90, CANX

# **Target/Specificity**

Calnexin

#### **Formulation**

 $50~\mu g$  (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

### **Reconstitution & Storage**

-20 °C

#### **Background Descriptions**

#### **Precautions**

Calnexin Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

## **Calnexin Blocking Peptide - Protein Information**

## Name CANX

#### **Function**

Calcium-binding protein that interacts with newly synthesized monoglucosylated glycoproteins in the endoplasmic reticulum. It may act in assisting protein assembly and/or in the retention within the ER of unassembled protein subunits. It seems to play a major role in the quality control apparatus of the ER by the retention of incorrectly folded proteins. Associated with partial T-cell



Tel: 858.875.1900 Fax: 858.875.1999



antigen receptor complexes that escape the ER of immature thymocytes, it may function as a signaling complex regulating thymocyte maturation. Additionally it may play a role in receptor-mediated endocytosis at the synapse.

#### **Cellular Location**

Endoplasmic reticulum membrane; Single-pass type I membrane protein. Mitochondrion membrane {ECO:0000250|UniProtKB:P24643}; Single-pass type I membrane protein. Melanosome membrane; Single-pass type I membrane protein. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:12643545, PubMed:17081065). The palmitoylated form preferentially localizes to the perinuclear rough ER (PubMed:22314232) Localizes to endoplasmic reticulum mitochondria-associated membrane (MAMs) that connect the endoplasmic reticulum and the mitochondria (By similarity). {ECO:0000250|UniProtKB:P24643, ECO:0000269|PubMed:12643545, ECO:0000269|PubMed:17081065, ECO:0000269|PubMed:22314232}

## Calnexin Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Calnexin Blocking Peptide - Images