

DGAT1 Blocking Peptide
Catalog # PBV10390b**Specification**

DGAT1 Blocking Peptide - Product Information

Primary Accession	O75907
Other Accession	EAW82128
Gene ID	8694
Calculated MW	55278

DGAT1 Blocking Peptide - Additional Information**Gene ID** 8694**Application & Usage**

The peptide is used for blocking the antibody activity of DGAT-1. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.

Other Names

Diacylglycerol O-acyltransferase 1, 2.3.1.20, ACAT-related gene product 1, Acyl-CoA retinol O-fatty-acyltransferase, ARAT, Retinol O-fatty-acyltransferase, 2.3.1.76, Diglyceride acyltransferase, DGAT1, AGRP1, DGAT

Target/Specificity

DGAT1

Formulation

50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

DGAT1 Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

DGAT1 Blocking Peptide - Protein Information**Name** DGAT1 {ECO:0000303|PubMed:16214399, ECO:0000312|HGNC:HGNC:2843}**Function**

Catalyzes the terminal and only committed step in triacylglycerol synthesis by using diacylglycerol

and fatty acyl CoA as substrates (PubMed:16214399, PubMed:18768481, PubMed:28420705, PubMed:9756920, PubMed:32433611, PubMed:32433610). Highly expressed in epithelial cells of the small intestine and its activity is essential for the absorption of dietary fats (PubMed:18768481). In liver, plays a role in esterifying exogenous fatty acids to glycerol, and is required to synthesize fat for storage (PubMed:16214399). Also present in female mammary glands, where it produces fat in the milk (By similarity). May be involved in VLDL (very low density lipoprotein) assembly (PubMed:18768481). In contrast to DGAT2 it is not essential for survival (By similarity). Functions as the major acyl-CoA retinol acyltransferase (ARAT) in the skin, where it acts to maintain retinoid homeostasis and prevent retinoid toxicity leading to skin and hair disorders (PubMed:16214399). Exhibits additional acyltransferase activities, including acyl CoA:monoacylglycerol acyltransferase (MGAT), wax monoester and wax diester synthases (By similarity). Also able to use 1-monoalkylglycerol (1-MAkG) as an acyl acceptor for the synthesis of monoalkyl-monoacylglycerol (MAMAG) (PubMed:28420705).

Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q9Z2A7}; Multi-pass membrane protein

DGAT1 Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DGAT1 Blocking Peptide - Images