10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# BLAME Blocking Peptide
**Catalog # PBV10420b**

## Specification

### BLAME Blocking Peptide - Product Information

| | |
|---|---|
| Primary Accession | D3ZS24 |
| Other Accession | NP_001099443 |
| Gene ID | 289237 |

### BLAME Blocking Peptide - Additional Information

| | |
|---|---|
| Application & Usage | **The peptide is used for blocking the antibody activity of BLAME. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.** |

**Target/Specificity**
BLAME

**Formulation**
50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

**Reconstitution & Storage**
-20 °C

**Background Descriptions**

**Precautions**
BLAME Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

### BLAME Blocking Peptide - Protein Information

### BLAME Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation

- Flow Cytomety
- Cell Culture

**BLAME Blocking Peptide - Images**