

PRMT-5 Blocking Peptide
Catalog # PBV10447b**Specification**

PRMT-5 Blocking Peptide - Product Information

Other Accession [EDM14175.1](#)
Gene ID **364382**

PRMT-5 Blocking Peptide - Additional Information**Application & Usage**

The peptide is used for blocking the antibody activity of PRMT-5. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.

Target/Specificity
PRMT-5

Formulation
50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

Reconstitution & Storage
-20 °C

Background Descriptions**Precautions**

PRMT-5 Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

PRMT-5 Blocking Peptide - Protein Information**PRMT-5 Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

PRMT-5 Blocking Peptide - Images