

CBP-Tag Blocking Peptide
Catalog # PBV10472b**Specification**

CBP-Tag Blocking Peptide - Product Information**CBP-Tag Blocking Peptide - Additional Information****Application & Usage**

The peptide is used for blocking the antibody activity of CBP-Tag. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.

Target/Specificity

CBP-tag

Formulation

50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

CBP-Tag Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

CBP-Tag Blocking Peptide - Protein Information**CBP-Tag Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CBP-Tag Blocking Peptide - Images

