

Human CellExp IL-10, murine recombinant protein
CSIF, IL-10, IL10A, TGIF, B-TCGF, GVHDS, MGC126450, MGC126451, RP11-262N9.1,
Interleukin-10
Catalog # PBV10895r

Specification

Human CellExp IL-10, murine recombinant protein - Product info

Primary Accession	P18893
Calculated MW	Calculated MW of 19.2 kDa with no tag. The predicted N-terminus is Ser 19. DTT-reduced protein migrates as 22.0 kDa due to glycosylation. KDa

Human CellExp IL-10, murine recombinant protein - Additional Info

Gene ID	16153
Gene Symbol	IL-10
Other Names	
CSIF, IL-10, IL10A, TGIF, B-TCGF, GVHDS, MGC126450, MGC126451, RP11-262N9.1, Interleukin-10	
Gene Source	Mouse
Source	HEK 293 cells
Assay&Purity	SDS-PAGE; ≥95%
Assay2&Purity2	N/A;
Recombinant	Yes
Target/Specificity	
IL-10	

Application Notes

Centrifuge the vial prior to opening. Reconstitute in sterile PBS, pH 7.4 to a concentration of 100 µg/ml. Do not vortex. This solution can be stored at 2-8°C for up to 1 month. For extended storage, it is recommended to store at -20°C.

Format

Lyophilized powder

Storage

-20°C; Lyophilized from 0.22 µm filtered solution in 20 mM Tris, 100 mM NaCl, pH 8.0. Generally 5-8% Mannitol or trehalose is added as a protectant before lyophilization.

Human CellExp IL-10, murine recombinant protein - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Human CellExp IL-10, murine recombinant protein - Images

Human CellExp IL-10, murine recombinant protein - Background

Interleukin-10 (IL-10) is also known as human cytokine synthesis inhibitory factor (CSIF), is an anti-inflammatory cytokine. IL-10 is an immunosuppressive cytokine produced by a variety of mammalian cell types including macrophages, monocytes, T cells, B cells and keratinocytes. Mature human IL-10 shares 72% - 86% amino acid sequence identity with bovine, canine, equine, feline, mouse, ovine, porcine, and rat IL-10. Whereas human IL-10 is active on mouse cells, mouse IL-10 does not act on human cells. IL-10 is capable of inhibiting synthesis of pro-inflammatory cytokines such as IFN- γ , IL-2, IL-3, TNF α and GM-CSF made by cells such as macrophages and regulatory T-cells. It also displays a potent ability to suppress the antigen-presentation capacity of antigen presenting cells. However, it is also stimulatory towards certain T cells and mast cells and stimulates B cell maturation and antibody production. Knockout studies suggested the function of Interleukin-10 / IL-10 as an essential immunoregulator in the intestinal tract. Patients with Crohn's disease react favorably towards treatment with bacteria producing recombinant interleukin-10, showing the importance of interleukin-10 for counteracting excessive immunity in the human body

Human CellExp IL-10, murine recombinant protein - References

Moore K.W., et al. Science 248:1230-1234(1990).
Kim J.M., et al. J. Immunol. 148:3618-3623(1992).
Church D.M., et al. PLoS Biol. 7:E1000112-E1000112(2009).
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Windsor W.T., et al. Biochemistry 32:8807-8815(1993).