

**Human CellExp CD85J/LILRB1, human recombinant protein**  
**CD85J, LILRB1, CD85, ILT2, LIR1, MIR7**  
**Catalog # PBV11112r****Specification****Human CellExp CD85J/LILRB1, human recombinant protein - Product info**Primary Accession  
Calculated MW[Q8NHL6](#)

This protein is fused with Fc fragment of human IgG1 at the C-terminus, has a calculated MW of 73.8 kDa. The predicted N-terminus is Gly 24. DTT-reduced Protein migrates as 90-110 kDa due to glycosylation. KDa

**Human CellExp CD85J/LILRB1, human recombinant protein - Additional Info**Gene ID  
Gene Symbol  
**Other Names**  
CD85J, LILRB1, CD85, ILT2, LIR1, MIR7**10859**  
**LILRB1**Gene Source  
Source  
Assay&Purity  
Assay2&Purity2  
Recombinant  
Results

**Human**  
**HEK293 cells**  
**SDS-PAGE; ≥95%**  
**N/A;**  
**Yes**  
**Measured by its ability to support the adhesion of HSB2 human peripheral blood acute lymphoblastic leukemia cells. Immobilized rhCD85J / LILRB1 Fc Chimera at 10 µg/mL, 100 µL/well can support 60-80% HSB2 cell adhesion, when 100000 cells were added in each well of a 96-well plate.**

**Target/Specificity**  
CD85J/LILRB1**Application Notes**

Centrifuge the vial prior to opening. Reconstitute in sterile PBS, pH 7.4 to a concentration of 50 µg/ml. Do not vortex. This solution can be stored at 2-8°C for up to 1 month. For extended storage, it is recommended to store at -20°C.

**Format**  
Lyophilized**Storage**

-20°C; Lyophilized from 0.22 µm filtered solution in PBS, pH 7.4. Normally Mannitol or Trehalose is added as protectants before lyophilization.

## **Human CellExp CD85J/LILRB1, human recombinant protein - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Human CellExp CD85J/LILRB1, human recombinant protein - Images**

## **Human CellExp CD85J/LILRB1, human recombinant protein - Background**

CD85 antigen-like family member J (CD85J) also known as Leukocyte immunoglobulin-like receptor subfamily B member 1 (LILRB1 or LIR-1), Immunoglobulin-like transcript 2 (ILT-2), Monocyte/macrophage immunoglobulin-like receptor 7 (MIR7), which belongs to leukocyte immunoglobulin-like receptor (LIR) family. CD85J / LILRB1 Contains 4 Ig-like C2-type (immunoglobulin-like) domains. CD85J / LIR-1 is expressed predominantly on B-cells and monocytes. CD85J is receptor for class I MHC antigens and recognizes a broad spectrum of HLA-A, HLA-B, HLA-C and HLA-G alleles and is also receptor for H301/UL18, a human cytomegalovirus class I MHC homolog. Ligand binding results in inhibitory signals and down-regulation of the immune response. CD85J / LILRB1 interaction with HLA-B or HLA-E leads to inhibition of the signal triggered by FCER1A and inhibits serotonin release. CD85J / LILRB1 inhibits FCGR1A-mediated phosphorylation of cellular proteins and mobilization of intracellular calcium ions.

## **Human CellExp CD85J/LILRB1, human recombinant protein - References**

Wagtmann N.,et al.Curr. Biol. 7:615-618(1997).  
Cosman D.,et al.Immunity 7:273-282(1997).  
Liu W.R.,et al.Immunogenetics 51:659-669(2000).  
Jones D.C.,et al.Eur. J. Immunol. 39:3195-3206(2009).  
Davidson C.L.,et al.Hum. Immunol. 71:942-949(2010).