

DDR2, Active recombinant protein
DDR, discoidin domain receptor tyrosine kinase 2
Catalog # PBV11334r**Specification**

DDR2, Active recombinant protein - Product info

Primary Accession	O16832
Concentration	0.1
Calculated MW	70.0 kDa KDa

DDR2, Active recombinant protein - Additional Info

Gene ID	4921
Gene Symbol	DDR2

Other Names

DDR, discoidin domain receptor tyrosine kinase 2, CD167 antigen-like family member B, Discoidin domain-containing receptor tyrosine kinase 2, Neurotrophic tyrosine kinase, receptor-related 3, Receptor protein-tyrosine kinase TKT, Tyrosine-protein kinase TYRO10

Source	Baculovirus (Sf9 insect cells)
Assay&Purity	SDS-PAGE; ≥90%
Assay2&Purity2	HPLC;
Recombinant	Yes
Format	
Liquid	

Storage

-80°C; Recombinant protein stored in 50mM Tris-HCl, pH 7.5, 150mM NaCl, 0.25mM DTT, 0.1mM EGTA, 0.1mM EDTA, 0.1mM PMSF, 25% glycerol.

DDR2, Active recombinant protein - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DDR2, Active recombinant protein - Images**DDR2, Active recombinant protein - Background**

DDR2 is a member of a novel subclass of RTKs containing a distinct extracellular region encompassing a factor VIII-like domain (1) and is thought to be involved in the regulation of cell

growth, differentiation, and metabolism. DDR2 plays a role in the regulation of collagen turnover mediated by smooth muscle cells in obstructive diseases of blood vessels and the lung (2).