

Human CellExp™ LYVE-1, mouse recombinant
Lymphatic Vessel Endothelial Hyaluronan (HA) Receptor-1, Xlkd1, Lyve-1, Crsbp-1
Catalog # PBV11494r

Specification

Human CellExp™ LYVE-1, mouse recombinant - Product info

Primary Accession [Q8BHC0](#)
Calculated MW **70 kDa** **KDa**

Human CellExp™ LYVE-1, mouse recombinant - Additional Info

Other Names

Lymphatic Vessel Endothelial Hyaluronan (HA) Receptor-1, Xlkd1, Lyve-1, Crsbp-1

Gene Source	Mouse
Source	HEK 293 cells
Assay&Purity	SDS-PAGE; ≥ 98%
Recombinant	Yes
Target/Specificity	
Lyve1	

Application Notes

Reconstitute in 1X PBS to the desired protein concentration.

Format

Lyophilized

Storage

-20°C; Lyophilized from 0.2 µm-filtered solution in PBS.

Human CellExp™ LYVE-1, mouse recombinant - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Human CellExp™ LYVE-1, mouse recombinant - Images

Human CellExp™ LYVE-1, mouse recombinant - Background

Lymphatic Vessel Endothelial Hyaluronan (HA) Receptor-1 (LYVE-1) is a 60-kDa type I transmembrane glycoprotein that is a member of the Link Protein superfamily. HA is found in the

extracellular matrix of most animal tissues and in body fluids. It modulates cell behavior and functions during tissue remodeling, development, homeostasis, and disease. It is often used as a marker of lymphatic endothelia. LYVE-1 is expressed on both the luminal and abluminal surfaces of lymphatic endothelium, and also on hepatic blood sinusoidal endothelia. This expression pattern, combined with studies showing that LYVE-1 can support cellular HA internalization in vitro, may suggest LYVE-1 participation in HA internalization for degradation, or transport of HA from tissues into the lumen of lymphatic vessels. LYVE-1-directed HA localization to lymphatic surfaces might also affect aspects of the immune response or tumor metastases. HA binding to CD44 can still occur in the presence of LYVE-1 in vitro. Therefore, LYVE-1-directed HA localization to lymphatics could provide a substrate for transmigrating CD44+ leukocytes or tumor cells. In addition to hepatic and lymphatic endothelia, some expression of LYVE-1 has been reported on Kupffer cells, the islets of Langerhans, cortical neurons, and renal epithelium.