

#### Crabrolin

Synthetic Peptide Catalog # SP2221a

## **Specification**

## **Crabrolin - Product Information**

Primary Accession Sequence

P01518

NH2-FLPLILRKIVTAL-CONH2

### **Crabrolin - Additional Information**

# **Other Names**

Crabrolin, CRBL

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

# Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## **Crabrolin - Protein Information**

## **Name CRBL**

## **Function**

Antimicrobial and mast cell degranulating peptide (PubMed:<a

href="http://www.uniprot.org/citations/6206053" target="\_blank">6206053</a>, PubMed:<a href="http://www.uniprot.org/citations/9273892" target="\_blank">9273892</a>, PubMed:<a href="http://www.uniprot.org/citations/31487493" target="\_blank">31487493</a>). Shows low antimicrobial activity towards some Gram-negative bacteria (E.coli MIC=300 ug/ml, S.typhimurium MIC=300 ug/ml) and all Gram-positive bacteria tested (B.subtilis MIC=300 ug/ml, S.aureus MIC=300 ug/ml, E.faecalis MIC=150 ug/ml, and E.faecium MIC=75 ug/ml) (PubMed:<a href="http://www.uniprot.org/citations/9273892" target="\_blank">9273892</a>, PubMed:<a href="http://www.uniprot.org/citations/31487493" target="\_blank">31487493</a>). May act by permeabilizing membranes, since it permeabilizes liposomes composed by phosphatidylglycerol(PG) and cardiolinin(CL)/phosphatidylglycerol(PG), which mimic the membranes of Gram-negative and

cardiolipin(CL)/phosphatidylglycerol(PG), which mimic the membranes of Gram-negative and Gram-positive bacteria, respectively (PubMed:<a

href="http://www.uniprot.org/citations/31487493" target="\_blank">31487493</a>). Causes hemolysis on rat and guinea pig erythrocytes (PubMed:<a

href="http://www.uniprot.org/citations/6206053" target="\_blank">6206053</a>, PubMed:<a href="http://www.uniprot.org/citations/9273892" target="\_blank">9273892</a>). Its mast cell degranulation activity may be related to the activation of G-protein coupled receptors in mast cells as well as interaction with other proteins located in cell endosomal membranes in the mast cells





(By similarity).

# **Cellular Location**

Secreted. Target cell membrane. Note=Assumes an amphipathic alpha-helical conformation in a membrane-like environment

# **Tissue Location**

Expressed by the venom gland.

**Crabrolin - Images**