

MUC5AC, Analog 1

Synthetic Peptide Catalog # SP2266b

### Specification

## MUC5AC, Analog 1 - Product Information

Primary Accession Sequence P98088 NH2-GTTPSPVPTTSTTSAP-COOH

### MUC5AC, Analog 1 - Additional Information

Gene ID 4586

**Other Names** Mucin-5AC, MUC-5AC, Gastric mucin, Lewis B blood group antigen, LeB, Major airway glycoprotein, Mucin-5 subtype AC, tracheobronchial, Tracheobronchial mucin, TBM, MUC5AC, MUC5

#### Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions** 

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### MUC5AC, Analog 1 - Protein Information

### Name MUC5AC {ECO:0000303|PubMed:11535137, ECO:0000312|HGNC:HGNC:7515}

Function

Gel-forming glycoprotein of gastric and respiratory tract epithelia that protects the mucosa from infection and chemical damage by binding to inhaled microorganisms and particles that are subsequently removed by the mucociliary system (PubMed:<a

href="http://www.uniprot.org/citations/14535999" target="\_blank">14535999</a>, PubMed:<a href="http://www.uniprot.org/citations/14718370" target="\_blank">14718370</a>). Interacts with H.pylori in the gastric epithelium, Barrett's esophagus as well as in gastric metaplasia of the duodenum (GMD) (PubMed:<a href="http://www.uniprot.org/citations/14535999" target="\_blank">14535999</a>).

Cellular Location Secreted

#### **Tissue Location**

Highly expressed in surface mucosal cells of respiratory tract and stomach epithelia. Overexpressed in a number of carcinomas. Also expressed in Barrett's esophagus epithelium and in the proximal duodenum.



# MUC5AC, Analog 1 - Images