10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Ubiquitin (65 - 76), (Ub2)

Synthetic Peptide
Catalog # SP2427b

Specification

Ubiquitin (65 - 76), (Ub2) - Product Information

Primary Accession	P68197
Other Accession	P69310, P19848, P62975, Q865C5, P69317
Sequence	**NH2-STLHLVLRLRGG-COOH**

Ubiquitin (65 - 76), (Ub2) - Additional Information

Other Names
Ubiquitin, UBIQ

Format
Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions
This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Ubiquitin (65 - 76), (Ub2) - Protein Information

Name UBIQ

Function
Ubiquitin exists either covalently attached to another protein, or free (unanchored). When covalently bound, it is conjugated to target proteins via an isopeptide bond either as a monomer (monoubiquitin), a polymer linked via different Lys residues of the ubiquitin (polyubiquitin chains) or a linear polymer linked via the initiator Met of the ubiquitin (linear polyubiquitin chains). Polyubiquitin chains, when attached to a target protein, have different functions depending on the Lys residue of the ubiquitin that is linked: Lys-48-linked is involved in protein degradation via the proteasome. Linear polymer chains formed via attachment by the initiator Met lead to cell signaling. Ubiquitin is usually conjugated to Lys residues of target proteins, however, in rare cases, conjugation to Cys or Ser residues has been observed. When polyubiquitin is free (unanchored-polyubiquitin), it also has distinct roles, such as in activation of protein kinases, and in signaling (By similarity).

Cellular Location
Cytoplasm. Nucleus.

Ubiquitin (65 - 76), (Ub2) - Images