

Oxyntomodulin / Glucagon 37 Synthetic Peptide Catalog # SP2842a

## **Specification**

# **Oxyntomodulin / Glucagon 37 - Product Information**

Primary Accession Other Accession Sequence

<u>08MJ25</u> <u>P01272</u>, <u>P01274</u>, <u>P29794</u> NH2-HSQGTFTSDYSKYLDSRRAQDFVQWLMN TKRNKNNIA-COOH

## **Oxyntomodulin / Glucagon 37 - Additional Information**

### **Other Names**

Glucagon, Glicentin, Glicentin-related polypeptide, GRPP, Oxyntomodulin, OXM, OXY, Glucagon, Glucagon-like peptide 1, GLP-1, Glucagon-like peptide 1(7-37), GLP-1(7-37), Glucagon-like peptide 1(7-36), GLP-1(7-36), Glucagon-like peptide 2, GLP-2, GCG

## Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

# **Oxyntomodulin / Glucagon 37 - Protein Information**

Name GCG

## Function

[Glucagon]: Plays a key role in glucose metabolism and homeostasis. Regulates blood glucose by increasing gluconeogenesis and decreasing glycolysis. A counterregulatory hormone of insulin, raises plasma glucose levels in response to insulin-induced hypoglycemia. Plays an important role in initiating and maintaining hyperglycemic conditions in diabetes.

## **Cellular Location**

Secreted {ECO:0000250|UniProtKB:P01275}.

### **Tissue Location**

Glucagon is secreted in the A cells of the islets of Langerhans. GLP-1, GLP-2, oxyntomodulin and glicentin are secreted from enteroendocrine cells throughout the gastrointestinal tract. GLP-1 and GLP-2 are also secreted in selected neurons in the brain

## Oxyntomodulin / Glucagon 37 - Images