

**ATG5 Antibody (N-term)**  
Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP1812a

**Specification**

**ATG5 Antibody (N-term) - Product Information**

|                      |  |
|----------------------|--|
| Application          | WB, IF, IHC-P,E  |
| Primary Accession    | <a href="#">Q9H1Y0</a>   |
| Other Accession      | <a href="#">Q3MQ06</a> , <a href="#">Q3MQ04</a> ,<br><a href="#">Q99J83</a> , <a href="#">Q3MQ24</a> ,<br><a href="#">Q6DEM4</a> |
| Reactivity Predicted | Human<br>Zebrafish, Bovine,<br>Mouse, Pig, Rat   |
| Host                 | Rabbit   |
| Clonality            | Polyclonal   |
| Isotype              | Rabbit Ig  |
| Antigen Region       | 1-30   |

**ATG5 Antibody (N-term) - Additional Information**

Gene ID 9474

**Other Names**

Autophagy protein 5, APG5-like,  
Apoptosis-specific protein, ATG5, APG5L, ASP

**Target/Specificity**

This ATG5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human ATG5.

**Dilution**

WB~~1:1000  
IF~~1:200  
IHC-P~~1:50~100

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ATG5 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.



All lanes : Anti-hAPG5L-D3 at 1:1000 dilution  
Lane 1: SH-SY5Y whole cell lysate Lane 2: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 32 kDa Blocking/Dilution buffer: 5% NFD/MTBST.



Fluorescent image of U251 cells stained with ATG5 (N-term) antibody. U251 cells were treated with Chloroquine (50 µM,16h), then fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.2%, 30 min). Cells were then incubated with AP1812a ATG5 (N-term) primary antibody (1:200, 2 h at room temperature). For secondary antibody, Alexa Fluor® 488

## ATG5 Antibody (N-term) - Protein Information

**Name** ATG5

**Synonyms** APG5L, ASP

### Function

Involved in autophagic vesicle formation. Conjugation with ATG12, through a ubiquitin-like conjugating system involving ATG7 as an E1-like activating enzyme and ATG10 as an E2-like conjugating enzyme, is essential for its function. The ATG12-ATG5 conjugate acts as an E3-like enzyme which is required for lipidation of ATG8 family proteins and their association to the vesicle membranes. Involved in mitochondrial quality control after oxidative damage, and in subsequent cellular longevity. Plays a critical role in multiple aspects of lymphocyte development and is essential for both B and T lymphocyte survival and proliferation. Required for optimal processing and presentation of antigens for MHC II. Involved in the maintenance of axon morphology and membrane structures, as well as in normal adipocyte differentiation. Promotes primary ciliogenesis through removal of OFD1 from centriolar satellites and degradation of IFT20 via the autophagic pathway.

### Cellular Location

Cytoplasm. Preautophagosomal structure membrane; Peripheral membrane protein  
Note=Colocalizes with nonmuscle actin. The conjugate detaches from the membrane immediately before or after autophagosome formation is completed (By similarity). Localizes also to discrete punctae along the ciliary axoneme and to the base of the ciliary axoneme

### Tissue Location


Ubiquitous. The mRNA is present at similar levels in viable and apoptotic cells, whereas the protein is dramatically highly expressed in apoptotic cells

## ATG5 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

conjugated donkey anti-rabbit antibody (green) was used (1:1000, 1h). Nuclei were counterstained with Hoechst 33342 (blue) (10 µg/ml, 5 min). ATG5 immunoreactivity is localized to autophagic vacuoles in the cytoplasm of U251 cells.



Formalin-fixed and paraffin-embedded human breast carcinoma tissue reacted with Autophagy APG5L antibody (N-term) (Cat.#AP1812a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

## ATG5 Antibody (N-term) - Background

Macroautophagy is the major inducible pathway for the general turnover of cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic enzymes and organelles during nutrient starvation. Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole). APG5, required for autophagy, conjugates to ATG12 and associates with an isolation membrane to form a cup-shaped isolation membrane and autophagosome. The conjugate detaches from the membrane immediately before or after autophagosome formation is completed. APG5 may also play an important role in the apoptotic process, possibly within the modified cytoskeleton. Its expression is a relatively late event in the apoptotic process, occurring downstream of caspase activity.

## ATG5 Antibody (N-term) - References

References for protein:

- 1.Baehrecke EH. Nat Rev Mol Cell Biol. 6(6):505-10. (2005)
2. Lum JJ, et al. Nat Rev Mol Cell Biol. 6(6):439-48. (2005)
- 3.Greenberg JT. Dev Cell. 8(6):799-801. (2005)
- 4.Levine B. Cell. 120(2):159-62. (2005)

5. Shintani T and Klionsky DJ. Science. 306(5698):990-5. (2004)
  6. Hammond E.M., et al. FEBS Lett. 425:391-395(1998)
  7. Strausberg R.L., et al. PNAS 99:16899-16903(2002)
  8. Grand R.J.A., et al. Exp. Cell Res. 218:439-451(1995)
  9. Mizushima N., et al. J. Biol. Chem. 273:33889-33892(1998)
  10. Mizushima N., et al. J. Cell Biol. 152:657-668(2001)
- References for U251 cell line:
1. Westermark B.; Pontén J.; Hugosson R. (1973). "Determinants for the establishment of permanent tissue culture lines from human gliomas". Acta Pathol Microbiol Scand A. 81:791-805. [PMID: 4359449].
  2. Pontén, J., Westermark B. (1978). "Properties of Human Malignant Glioma Cells in Vitro". Medical Biology 56: 184-193.[PMID: 359950].
  3. Geng Y.; Kohli L.; Klocke B.J.; Roth K.A.(2010). "Chloroquine-induced autophagic vacuole accumulation and cell death in glioma cells is p53 independent". Neuro Oncol. 12(5): 473-481.[PMID: 20406898].

### ATG5 Antibody (N-term) - Citations

- [ATG5 Promotes Death Signaling in Response to the Cyclic Depsipeptides Coibamide A and Apratoxin A.](#)
- [Atg5flox-Derived Autophagy-Deficient Model of Pompe Disease: Does It Tell the Whole Story?](#)
- [The Detection Techniques for Autophagy-Associated Cell Death-Related Genes and Proteins: Gene Expression Assay and Immunohistochemistry.](#)
- [Effects of Combined Lysosomal and Mitochondrial Photodamage in a Non Small-Cell Lung Cancer Cell Line: the Role of Paraptosis.](#)
- [Studying Autophagy in Zebrafish.](#)
- [Promotion of Pro-Apoptotic Signals by Lysosomal Photodamage: Mechanistic Aspects and Influence of Autophagy.](#)
- [FGFR3/Fibroblast Growth Factor Receptor 3 Inhibits Autophagy through Decreasing the ATG12-ATG5 Conjugate, Leading to the Delay of Cartilage Development in Achondroplasia.](#)
- [Autophagy in spinal motor neurons of conditional ADAR2-knockout mice: an implication for a role of calcium in increased autophagy flux in ALS.](#)
- [Long-term artificial selection reveals a role of TCTP in autophagy in mammalian cells.](#)
- [Coxsackievirus B3 induces crosstalk between autophagy and apoptosis to benefit its release after replicating in autophagosomes through a mechanism involving caspase cleavage of autophagy-related proteins.](#)
- [Control of photoreceptor autophagy after retinal detachment: the switch from survival to death.](#)
- [Autophagosomes contribute to intracellular lipid distribution in enterocytes.](#)
- [Autophagy Induced by Tumor Necrosis Factor  \$\alpha\$  Mediates Intrinsic Apoptosis in Trophoblastic Cells.](#)
- [Down-regulation of autophagy-related protein 5 \(ATG5\) contributes to the pathogenesis of early-stage cutaneous melanoma.](#)
- [Phosphorylation of Atg5 by the Gadd45<sup>1</sup>-MEKK4-p38 pathway inhibits autophagy.](#)
- [Cell loss and autophagy in the extra-adrenal chromaffin organ of Zuckerkandl are regulated by glucocorticoid signalling.](#)
- [Arsenic trioxide enhances the radiation sensitivity of androgen-dependent and -independent human prostate cancer cells.](#)
- [Autophagy induced by deficiency of sphingosine-1-phosphate phosphohydrolase 1 is switched to apoptosis by calpain-mediated autophagy-related gene 5 \(Atg5\) cleavage.](#)
- [Expression pattern and functions of autophagy-related gene atg5 in zebrafish organogenesis.](#)
- [Arsenic trioxide induces autophagy and apoptosis in human glioma cells in vitro and in vivo through downregulation of survivin.](#)
- [A rapid method to improve protein detection by indirect ELISA.](#)
- [p62, Ref\(2\)P and ubiquitinated proteins are conserved markers of neuronal aging, aggregate formation and progressive autophagic defects.](#)
- [Sphingosine-1-phosphate phosphohydrolase-1 regulates ER stress-induced autophagy.](#)
- [Control of basal autophagy by calpain1 mediated cleavage of ATG5.](#)

- An increase in intracellular  $\text{Ca}^{2+}$  is required for the activation of mitochondrial calpain to release AIF during cell death.